## EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION ORGANISATION EUROPEENNE ET MEDITERRANEENNE POUR LA PROTECTION DES PLANTES

(11-17239)

## Summary sheet of validation data for a diagnostic test

The EPPO Standard PM 7/98 *Specific requirements for laboratories preparing accreditation for a plant pest diagnostic activity* describes how validation should be conducted. It also includes definitions of performance criteria.

Target Organism	Acidovorax citrulli			
Short description	Detection of Acidovorax citrulli by PCR in seeds			
Laboratory contact details	Naktuinbouw Sotaweg 22, 2371 GD Roelofarendsveen, Netherlands			
Date and reference of the validation report	2015-11-16 - v1.2			
Validation process according to EPPO Standard PM 7/98:	Yes			
Reference of the test description	N/R The protocol will be included in the EPPO diagnostic protocol that is currently being drafted.			
Is the test the same as described in the EPPO DP?				
Is the lab accredited for this test?	No			
Plant species tested (if relevant)	Citrullus lanatus, Cucumis melo and other Cucurbitaceae			
Matrices tested (if relevant)	seeds and isolates			
List of methods used				
Method for extraction / isolation / baiting of target organism from matrix				
Molecular methods, e.g. hybridization, PCR and real time PCR	х	DNA extraction using Kingfisher and Sbeadex maxi kit for Acidovorax citrulli (LGC Genomics) followed by Taqman PCR		
Serological methods: IF, ELISA, Direct Tissue Blot Immuno Assay				
Plating methods: selective isolation				
•	_	-		

Bioassay methods: selective enrichment in host plants, baiting, plant test and grafting.				
Pathogenicity test				
Fingerprint methods: protein profiling, fatty acid profiling & DNA profiling				
Morphological and morphometrical methods intended for identification				
Biochemical methods: e.g. enzyme electrophoresis, protein profiling				
Other				
Analytical sensitivity (= limit of detect	<u>ion)</u>			
What is smallest amount of target that can be detected reliably?	The limit of detection at 0.95 probability is 9 cells/mL seed extract.			
<u>Diagnostic sensitivity</u>				
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98	Not determined			
Specify the standard test	No standard test available			
<u>Analytical specificity</u>				
Specificity value	Diagnostic sensitivity: 100% Diagnostic specificity: 98%			
Number of strains/populations of target organisms tested	168 strains			
Number of non-target organisms tested	54 non-targets			
Cross reacts with (specify the species)	Two primers sets tested - IS1002 cross-reacts with 2 of the 9 Acidovorax cattleyae isolates tested and several unknown bacteria - Contig22 cross-reacts only with 1 unknown bacteria, characterized in AFLP-study, outside of the Acit tree. Cross-reacts with both primer sets			
Diagnostic Specificity				
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test	98%			
Specify the standard test	AFLP-study			

<b> </b>	•		
Reproducibility			
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	100% Intralaboratory testing		
Repeatability			
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	100% Inter- and intralaboratory testing		
Test performance study			
Test performance study?	No		
Include brief details of the test performance study and its output.It available, provide a link to published article/report			
Other information			
Any other information considered useful e.g. robustness, ease of performing the test, etc.			
The following complementary files are available online:	<ul> <li>Poster Koenraadt et al 2014 ISTA         Seed Health Symposium</li> <li>Specificity of Contig21 Taqman</li> <li>Validation report Acidovorax citrulli v1.2</li> </ul>		