EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION ORGANISATION EUROPEENNE ET MEDITERRANEENNE POUR LA PROTECTION DES PLANTES

(11-17239)

Summary sheet of validation data for a diagnostic test

The EPPO Standard PM 7/98 *Specific requirements for laboratories preparing accreditation for a plant pest diagnostic activity* describes how validation should be conducted. It also includes definitions of performance criteria.

	Durcarl	a alan chuic yydan hilus	
Target Organism	Bursaphelenchus xylophilus		
Short description	Duplex PCR based identification of Bursaphelenchus xylophilus Jiang et al. 2005		
Laboratory contact details	Anses, Laboratoire de la Santé des Végétaux - Unité de Nématologie Domaine de la Motte au Viconte BP 35327, 35653 Le Rheu, France		
Date and reference of the validation report	2011-05 - Evaluation d'outils moléculaires d'identification de Bursaphelenchus xylophilus sur individus isolés		
Validation process according to EPPO Standard PM 7/98:	Yes		
Reference of the test description	0 Jiang L, Zheng J, Waeyenberge L, Subbotin S, Moens M (2005) Duplex PCR based identification of Bursaphelenchus xylophilus (Steiner & Buhrer,1934) Nickle, 1970. Russian Journal of Nematology, 2005, 13(2), 115-121		
Is the test the same as described in the EPPO DP?	No Cited but not described in full		
Is the lab accredited for this test?	No		
Plant species tested (if relevant)			
Matrices tested (if relevant)	Nematodes		
List of methods used			
Method for extraction / isolation / baiting of target organism from matrix			
Molecular methods, e.g. hybridization, PCR and real time PCR	Х	Two steps PCR : 1 universal PCR + 1 species specific PCR	
Serological methods: IF, ELISA,			

Direct Tissue Blot Immuno Assay				
Plating methods: selective isolation				
Bioassay methods: selective enrichment in host plants, baiting, plant test and grafting.				
Pathogenicity test				
Fingerprint methods: protein profiling, fatty acid profiling & DNA profiling				
Morphological and morphometrical methods intended for identification				
Biochemical methods: e.g. enzyme electrophoresis, protein profiling				
Other				
Analytical sensitivity (= limit of detection)				
What is smallest amount of target that can be detected reliably?	5 nematodes			
<u>Diagnostic sensitivity</u>				
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98	not performed			
Specify the standard test	not relevant			
<u>Analytical specificity</u>				
Specificity value	100%			
Number of strains/populations of target organisms tested	7 populations (see table1)			
Number of non-target organisms tested				
Cross reacts with (specify the species)	15 populations (see table1)			
Diagnostic Specificity	Diagnostic Specificity			
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test	Not performed			
Specify the standard test	Not relevant			
<u>Reproducibility</u>				
Provide the calculated % of				
ı	I	·		

agreement for a given level of the pest (see PM 7/98)	100% for 5 individuals		
Repeatability			
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	100% for 5 individuals		
Test performance study			
Test performance study?	No		
Include brief details of the test performance study and its output.It available, provide a link to published article/report			
Other information			
Any other information considered useful e.g. robustness, ease of performing the test, etc.	see table 2. The full report is available upon request to the laboratory.		
The following complementary files are available online:	 comparison of different PCR tests Table 1 Specificity_PCR 		