## EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION ORGANISATION EUROPEENNE ET MEDITERRANEENNE POUR LA PROTECTION DES PLANTES Summary sheet of validation data for a diagnostic test

The EPPO Standard PM 7/98 Specific requirements for laboratories preparing accreditation for a plant pest diagnostic activity describes how validation should be conducted. It also includes definitions of performance criteria.

Laboratory contact details	National Institiute of Biology, Department of Biotechnology and Systems Biology Vecna pot 121, 1000 Ljubljana, Slovenia
Short description of the test	Detection of Xylella fastidiosa by real-time PCR in plant material
Date, reference of the validation report	2016-02-26 - Dreo, Tanja, 2016. Validation data on the modified real-time PCR for detection of Xylella fastidiosa adapted from Francis et al. (2006) (No. D0002/16). National Institute of Biology, Department of Biotechnology and Systems Biology, Ljubljana.
Validation process according to EPPO Standard PM7/98?	no
Is the lab accredited for this test?	no
Was the validated data generated in the framework of a project?	
Description of the test	
Organism(s)	Xylella fastidiosa (XYLEFA)
Detection / identification	detection
Method(s)	Molecular Extraction DNA RNA Molecular real time PCR
Method: Molecular Extraction DNA RNA	
Reference of the test description	
As or adapted from an EPPO diagnostic protocol	yes
EPPO Diagnostic Protocol name	PM 7/024 Xylella fastidiosa (version 2)
As or adapted from an IPPC diagnostic protocol	no
Kit	
Is a kit used	yes
Manufacturer name	BIONOBILE
Specify the kit used	QuickPick™ SML Plant DNA
Kit used following the manufacturer's instructions?	

Other information		
Other details on the test	DNA extraction from plant material using QuickPick™ SML Plant DNA kit (Bionobile).	
Method: Molecular real time PCR		
Reference of the test description		
As or adapted from an EPPO diagnostic protocol	yes	
EPPO Diagnostic Protocol name	PM 7/024 Xylella fastidiosa (version 2)	
Name of the test	Taqman real-time PCR tests (based on Francis et al., 2006)	
As or adapted from an IPPC diagnostic protocol	no	
Other information		
Reaction type	Simplex - Probe	
Other details on the test	Modified real-time PCR adapted from Francis, M., Lin, H., Rosa, J.CL., Doddapaneni, H., Civerolo, E.L., 2006. Genome-based PCR Primers for Specific and Sensitive Detection and Quantification of Xylella fastidiosa. European Journal of Plant Pathology 115, 203–213. doi:10.1007/s10658-006-9009-4	
Are the performance characteristics included in the EPPO diagnostic protocol?	yes	
Performance Criteria :		
Organism 1.:	Xylella fastidiosa(XYLEFA)	
Analytical sensitivity		
What is smallest amount of target that can be detected reliably?	On the DNA extracted from pure cultures of X. fastidiosa: 2.6, 3.2 and 3.5 (log (cells/mL) of Xylella fastidiosa subsp. multiplex, Xylella fastidiosa, and Xylella fastidiosa subsp. pauca CoDiRO strain, respectively. On plant material: 94 % (determined on log 5 cells/mL of plant extracts; the lowest	
	concentration tested)	
Diagnostic sensitivity	· '	
Diagnostic sensitivity  Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98	· '	
Proportion of infected/infested samples tested positive compared to results from the	concentration tested)	
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98	concentration tested)	
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98  Analytical specificity - inclusivity  Number of strains/populations of target	No data available.	
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98  Analytical specificity - inclusivity  Number of strains/populations of target organisms tested	No data available.	
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98  Analytical specificity - inclusivity  Number of strains/populations of target organisms tested  Specificity value	No data available.	

Diagnostic Specificity		
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test	No data available.	
Reproducibility		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	0,97	
Repeatability		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	No data avilable.	
Test performance study		
Test performance study?	no	
·		
The following complementary files are available online:	<ul> <li>Validation data on the modified real-time PCR for detection of Xylella fastidiosa adapted from Francis et al. (2006) (No. D0002/16)</li> </ul>	

Creation date: 2016-02-26 00:00:00 - Last update: 2020-10-08 18:56:50