EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION ORGANISATION EUROPEENNE ET MEDITERRANEENNE POUR LA PROTECTION DES PLANTES Summary sheet of validation data for a diagnostic test

The EPPO Standard PM 7/98 Specific requirements for laboratories preparing accreditation for a plant pest diagnostic activity describes how validation should be conducted. It also includes definitions of performance criteria.

Laboratory contact details	Anses Plant Health Laboratory - Pests and Tropical Pathogens Unit Pôle de Protection des Plantes, 7 Chemin de l'IRAT, 97410 Saint Pierre, France
Short description of the test	Detection and identification of 'Candidatus Liberibacter asiaticus' and 'Candidatus Liberibacter africanus' by Molecular real time PCR (according to Bertolini et al. (2010, 2014)) in Citrus sp. leaves
Date, reference of the validation report	2020-07-10 - HLB_qPCR_EUPHRESCO-2016-A-232
Link to other validation data	- HLB_qPCR_EUPHRESCO-2016-A-232 Detection and identification of 'Candidatus Liberibacter asiaticus' and 'Candidatus Liberibacter africanus' by Molecular real time PCR (according to Bertolini et al. (2010, 2014)) in Citrus sp. leaves
Validation process according to EPPO Standard PM7/98?	yes
Is the lab accredited for this test?	no
Was the validated data generated in the framework of a project?	Euphresco
If yes, please specify	2016-A-232
Description of the test	
Organism(s)	'Candidatus Liberibacter africanus' (LIBEAF) 'Candidatus Liberibacter asiaticus' (LIBEAS)
Detection / identification	detection and identification
Method(s)	Molecular Extraction DNA RNA Molecular real time PCR
Method: Molecular Extraction DNA RNA	
Reference of the test description	
Kit	
Is a kit used	yes
Manufacturer name	QIAGEN
Specify the kit used	DNeasy Plant Mini Kit
Kit used following the manufacturer's instructions?	yes DNA extraction was performed on ground citrus leaves using the DNeasy Plant Mini Kit (Qiagen, 1 / 4 Germantown, MD, USA) following the

Ground using a HOMEX 6 homogenizer (Bioreba AG, Reinach, Switzerland) with 5 mL of extraction buffer (pH = 8): 50 mM Sigma 7-9® TRIS (Merck KGaA), Darmstadt, Germany); 5 mM EDTA (Merck KGaA), and 1%sodium dodecyl sulfate (Merck KGaA), and 1%sodium dodecyl sulfate (Merck KGaA), and 1%sodium dodecyl sulfate (Merck KGaA). Method: Molecular real time PCR Reference of the test description As or adapted from an EPPO diagnostic protocol EPPO Diagnostic Protocol name PM 7/121 < >(-		manufacturer's recommendations.	
AG. Reinach, Switzerland) with 5 ml, of extraction buffer (pH = 8): 50 mM Sigma 7-9® TRIS (Merck KGaA, Darmstadt, Germany): 5 mM EDTA (Merck KGaA); and 1%sodium dodecyl sulfate (Merck KGaA). Method: Molecular real time PCR Reference of the test description As or adapted from an EPPO diagnostic protocol EPPO Diagnostic Protocol name PM 7/121 <i>'Candidatus </i> ' Elberibacter africanus', ' <i <="" candidatus="" i="">' Elberibacter americanus' and ' <i <="" candidatus="" i="">' Elberibacter africanus', ' <i <="" candidatus="" i=""> Name of the test Real-time PCR targeting 16S rRNA gene (according to Bertolini et al. (2010, 2014)) Is the test modified compared to the reference test Kit Is a kit used Other information Reaction type Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1: 'Candidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - Inclusivity Number of strains/populations of target</i></i></i></i></i></i></i>	Other information		
Reference of the test description As or adapted from an EPPO diagnostic protocol EPPO Diagnostic Protocol name PM 7/121 < >'Candidatus 'Candidatus 'Candidatus</ Candidatus</ C</th <th>Other details on the test</th> <th>AG, Reinach, Switzerland) with 5 mL of extraction buffer (pH = 8): 50 mM Sigma 7-9® TRIS (Merck KGaA, Darmstadt, Germany); 5 mM EDTA (Merck KGaA); and 1%sodium dodecyl sulfate (Merck</th>	Other details on the test	AG, Reinach, Switzerland) with 5 mL of extraction buffer (pH = 8): 50 mM Sigma 7-9® TRIS (Merck KGaA, Darmstadt, Germany); 5 mM EDTA (Merck KGaA); and 1%sodium dodecyl sulfate (Merck	
As or adapted from an EPPO diagnostic protocol EPPO Diagnostic Protocol name PM 7/121 <i>'Candidatus</i> Liberibacter africanus', '< > Candidatus Liberibacter africanus', '< > Candidatus Liberibacter africanus', '< > Candidatus Liberibacter asiaticus' (version 1) Name of the test Real-time PCR targeting 16S rRNA gene (according to Bertolini et al. (2010, 2014)) Name of the test modified compared to the reference test Kit Is a kit used no Other information Reaction type Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: Candidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target Number of strains/populations of target	Method: Molecular real time PCR		
PPO Diagnostic Protocol name PM 7/121 <i>'Candidatus </i> Liberibacter africanus', 'Icandidatus Real-time PCR targeting 16S rRNA gene (according to Bertolini et al. (2010, 2014)) As or adapted from an IPPC diagnostic protocol Is the test modified compared to the reference test Kit Is a kit used In Italy is a kit used in Italy is a kit used In Italy is a kit used in Italy is a kit used Italy is a kit used in Italy is a kit used Italy is a kit used Italy is a kit used in Italy is a kit used in Italy is a kit used Italy is a kit used in Italy is a kit used in Italy is a kit used in Italy is a kit used Italy is a kit used in Italy is a kit used is a kit used in Italy is a kit used is a kit used in Italy is a kit used is a kit used in Italy is a	Reference of the test description		
africanus', ' <i>Candidatus</i> Liberibacter americanus' and ' <i>Candidatus</i> Liberibacter asiaticus' (version 1) Name of the test Real-time PCR targeting 165 rRNA gene (according to Bertolini et al. (2010, 2014)) As or adapted from an IPPC diagnostic protocol Is the test modified compared to the reference test Kit Is a kit used no Other information Reaction type Simplex - Probe Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: 'Candidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity. Number of strains/populations of target Number of strains/populations of target Real-time PCR targeting 165 rRNA gene (according to Senatics' (2010, 2014)) no 10 10 10 10 11 11 12 13 14 15 15 16 16 16 16 17 16 17 17 17 18 18 18 18 18 18 18	As or adapted from an EPPO diagnostic protocol	yes	
to Bertolini et al. (2010, 2014)) As or adapted from an IPPC diagnostic protocol Is the test modified compared to the reference test Kit Is a kit used Other information Reaction type Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: Candidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	EPPO Diagnostic Protocol name	africanus', ' <i>Candidatus</i> Liberibacter americanus' and ' <i>Candidatus</i> Liberibacter	
Is the test modified compared to the reference test Kit Is a kit used Other information Reaction type Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: Vandidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target Incompared to the properties of the proportion of the proportion of the properties of the proportion of the properties of the p	Name of the test		
reference test Kit Is a kit used no Other information Reaction type Simplex - Probe Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: 'Candidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	As or adapted from an IPPC diagnostic protocol	no	
Other information Reaction type Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target Simplex - Probe no (Candidatus Liberibacter africanus'(LIBEAF) (LIBEAF) (DSE=PA/N+, Fav. Hypothesis, considering CLas and CLaf samples)	Is the test modified compared to the reference test	no	
Cother information Reaction type Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target Simplex - Probe no (Candidatus Liberibacter africanus'(LIBEAF)	Kit		
Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target Simplex - Probe no Candidatus Liberibacter africanus'(LIBEAF) 'Candidatus Liberibacter africanus'(LIBEAF) 'Candidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity 1 (DSE=PA/N+, Fav. Hypothesis, considering CLas and CLaf samples) 1 (DSE=PA/N+, Fav. Hypothesis, considering CLas and CLaf samples) Candidatus Liberibacter africanus'(LIBEAF) Ca	Is a kit used	no	
Are the performance characteristics included in the EPPO diagnostic protocol? Performance Criteria: Organism 1.: Candidatus Liberibacter africanus'(LIBEAF) Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	Other information		
Performance Criteria: Organism 1.: Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target Candidatus Liberibacter africanus (LIBEAF) Candidatus Liberibacte	Reaction type	Simplex - Probe	
Organism 1.: Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target 'Candidatus Liberibacter africanus'(LIBEAF) 'Candidatus Liberibacter africanus'(LIBEAF) I (DSE=PA/N+, Fav. Hypothesis, considering CLas and CLaf samples)	Are the performance characteristics included in the EPPO diagnostic protocol?	no	
Analytical sensitivity What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	Performance Criteria :		
What is smallest amount of target that can be detected reliably? Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	Organism 1.:	'Candidatus Liberibacter africanus'(LIBEAF)	
Diagnostic sensitivity Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	Analytical sensitivity		
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	What is smallest amount of target that can be detected reliably?		
tested positive compared to results from the standard test, see appendix 2 of PM 7/98 Analytical specificity - inclusivity Number of strains/populations of target	Diagnostic sensitivity		
Number of strains/populations of target	Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98	· · · · · · · · · · · · · · · · · · ·	
· · · · · · · · · · · · · · · · · · ·	Analytical specificity - inclusivity		
urganisins testeu	Number of strains/populations of target organisms tested		
Specificity value	Specificity value		
Analytical specificity - exclusivity			
Number of non-target organisms tested	Number of non-target organisms tested		

Specificity value		
Diagnostic Specificity		
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test	0.449 (DSP=NA/N-, Fav. Hypothesis, considering CLas and CLaf samples)	
Reproducibility		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	CO=0.865 (considering CLas and CLaf samples)	
Repeatability		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	DA=0.988 (considering CLas and CLaf samples)	
Organism 2.:	'Candidatus Liberibacter asiaticus'(LIBEAS)	
Diagnostic sensitivity		
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98	1 (DSE=PA/N+, Fav. Hypothesis, considering CLas and CLaf samples)	
Diagnostic Specificity		
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test	0.449 (DSP=NA/N-, Fav. Hypothesis, considering CLas and CLaf samples)	
Reproducibility		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	CO=0.865 (considering CLas and CLaf samples)	
Repeatability		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	DA=0.988 (considering CLas and CLaf samples)	
Test performance study		
Test performance study?	yes	
Brief details of the test performance study and its output.It available, link to published article/report	Test performance study organized in the framework of a EUPHRESCO project involving 8 international laboratories.	
Other information		
Any other information considered useful	Publication available at: https://link.springer.com/content/pdf/10.1007/s10658-020-02052-3.pdf Cellier, G., C. Redondo, J. Cubero, M. Roselló, E. de Andrade, L. Cruz, E. Ince, H. N. Yildiz, P. G. Güler, A. M. D'Onghia, T. Yaseen, K. Djelouah, E. Metz-Verschure, F. Gaffuri, R. A. Gottsberger, and B. Giovani. 2020. "Comparison of the performance of the main real-time and conventional PCR detection tests for 'Candidatus Liberibacter' spp., plant pathogenic bacteria causing the Huanglongbing disease in Citrus spp." European Journal of Plant Pathology. doi: 10.1007/s10658-020-02052-3.	

Creation date: 2020-08-18 15:22:57 - Last update: 2020-09-14 15:24:40