## EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION ORGANISATION EUROPEENNE ET MEDITERRANEENNE POUR LA PROTECTION DES PLANTES (11-17239)

## Summary sheet of validation data for a diagnostic test

The EPPO Standard PM 7/98 Specific requirements for laboratories preparing accreditation for a plant pest diagnostic activity describes how validation should be conducted. It also includes definitions of performance criteria.

	T		
Target Organism	Globodera pallida Globodera rostochiensis		
Short description	Molecular identification of Globodera pallida and G. rostochiensis - PCR-RFLP		
Laboratory contact details	Anses Plant Health Laboratory - Nematology Unit Domaine de la Motte au Viconte BP 35327, 35653 Le Rheu, France		
Date and reference of the validation report	13/07/2010 - Evaluation comparative de différents outils moléculaire spour l'identfiication de Globodera pallida et G. rostochiensis - ref 10/03		
Validation process according to EPPO Standard PM 7/98:	Yes		
Reference of the test description	0 PM7/40 (2) APPENDIX 3 - PART C Thiéry & Mugniéry (1996) Interspecific rDNA restriction fragment length polymorphism in Globodera species, parasites of Solanaceous plants. Fundamental and Applied Nematology 19, 471–479.		
Is the test the same as described in the EPPO DP?	Yes		
Is the lab accredited for this test?	No		
Plant species tested (if relevant)			
Matrices tested (if relevant)	isolated nematodes		
List of methods used			
Method for extraction / isolation / baiting of target organism from matrix			
Molecular methods, e.g. hybridization, PCR and real time PCR	Х	PCR-RFLP	
Serological methods: IF, ELISA, Direct Tissue Blot Immuno Assay			
Plating methods: selective isolation			
Bioassay methods: selective enrichment in host plants, baiting, plant test and grafting.			
Pathogenicity test			

Fingerprint methods: protein profiling, fatty acid profiling & DNA profiling				
Morphological and morphometrical methods intended for identification				
Biochemical methods: e.g. enzyme electrophoresis, protein profiling				
Other				
Analytical sensitivity (= limit of detection)				
What is smallest amount of target that can be detected reliably?	1 individual of J2 stage			
Diagnostic sensitivity				
Proportion of infected/infested samples tested positive compared to results from the standard test, see appendix 2 of PM 7/98				
Specify the standard test				
Analytical specificity				
Specificity value		. pallida (1 non target species detected among 11) G. rostochiensis		
Number of strains/populations of target organisms tested	11 populations of G. pallida and 4 populations of G. rostochiensis (refer to evaluation report 10/03 - annex 2)			
Number of non-target organisms tested	7 populations tested (list of populations, refer to evaluation report 10/03 - annex 2)			
Cross reacts with (specify the species)	no cross reaction			
Diagnostic Specificity				
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test				
Specify the standard test				
Reproducibility				
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	90% for G. pallida (at the level of 1 J2 stage) 93% for G. rostochiensis (at the level of 1 J2 stage)			
Repeatability				
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	100% for both species			
Test performance study				
Test performance study?	No			
Include brief details of the test performance study and its output.It available, provide a link to				

published article/report	
Other information	
Any other information considered useful e.g. robustness, ease of performing the test, etc.	presence of unpublished profiles for South American populations of G. pallida. The full report is available upon request to the laboratory.