

EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION
ORGANISATION EUROPEENNE ET MEDITERRANEENNE POUR LA PROTECTION DES PLANTES
(11-17239)

Summary sheet of validation data for a diagnostic test

The EPPO Standard PM 7/98 *Specific requirements for laboratories preparing accreditation for a plant pest diagnostic activity* describes how validation should be conducted. It also includes definitions of performance criteria.

Target Organism	Meloïdogyne chitwoodi Meloïdogyne fallax	
Short description	Detection of Meloïdogyne chitwoodi and M. fallax by PCR RFLP	
Laboratory contact details	Anses, Laboratoire de la Santé des Végétaux - Unité de Nématologie Domaine de la Motte au Viconte BP 35327, 35653 Le Rheu, France	
Date and reference of the validation report	september 2010 - validation report - september 2010	
Validation process according to EPPO Standard PM 7/98:	Yes	
Reference of the test description	N/R Zijlstra et al. (1995). Differences between ITS regions of isolates of root-knot nematodes Meloïdogyne hapla and M. chitwoodi . Phytopathology 85, 1231-1237.	
Is the test the same as described in the EPPO DP?	No not included in PM7/41 (Zijlstra et al. 1997 included)	
Is the lab accredited for this test?	Yes	
Plant species tested (if relevant)		
Matrices tested (if relevant)	isolated nematodes	
List of methods used		
Method for extraction / isolation / baiting of target organism from matrix		
Molecular methods, e.g. hybridization, PCR and real time PCR	X	PCR RFLP
Serological methods: IF, ELISA, Direct Tissue Blot Immuno Assay		
Plating methods: selective isolation		
Bioassay methods: selective enrichment in host plants, baiting, plant test and grafting.		
Pathogenicity test		
Fingerprint methods: protein profiling, fatty acid profiling & DNA		

profiling		
Morphological and morphometrical methods intended for identification		
Biochemical methods: e.g. enzyme electrophoresis, protein profiling		
Other		
<u>Analytical sensitivity (= limit of detection)</u>		
What is smallest amount of target that can be detected reliably?	1 J2 for M. fallax and 1 J2 for M. chitwoodi	
<u>Diagnostic sensitivity</u>		
Proportion of infected/infested samples tested positive compared to results from the standard test , see appendix 2 of PM 7/98		
Specify the standard test		
<u>Analytical specificity</u>		
Specificity value	100% for M. fallax and 100% for M. chitwoodi	
Number of strains/populations of target organisms tested	1 population for M. fallax and 4 populations for M. chitwoodi (for details see annex 1 of validation report)	
Number of non-target organisms tested	29 nematodes populations (see Annex 1 of validation report)	
Cross reacts with (specify the species)	none	
<u>Diagnostic Specificity</u>		
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test		
Specify the standard test		
<u>Reproducibility</u>		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	87% for 1 J2 and 100% for 2 J2 for M. fallax; 66% for 1 J2 and 100% for 2 J2 of M. chitwoodi	
<u>Repeatability</u>		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	50% for 1 J2 and 100% for 2 J2 for M. fallax; 25% for 1 J2 and 100% for 2 J2 of M. chitwoodi	
<u>Test performance study</u>		
Test performance study?	No	
Include brief details of the test performance study and its output.It available, provide a link to published article/report		
<u>Other information</u>		

**Any other information considered useful
e.g. robustness, ease of performing the test, etc.**

The full report is available upon request to the laboratory.