

EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION
ORGANISATION EUROPEENNE ET MEDITERRANEENNE POUR LA PROTECTION DES PLANTES
(11-17239)

Summary sheet of validation data for a diagnostic test

The EPPO Standard PM 7/98 *Specific requirements for laboratories preparing accreditation for a plant pest diagnostic activity* describes how validation should be conducted. It also includes definitions of performance criteria.

Target Organism	Meloidogyne enterolobii	
Short description	Identification of meloidogyne enterolobii by SCAR PCR	
Laboratory contact details	Anses Plant Health Laboratory - Nematology Unit Domaine de la Motte au Viconte BP 35327, 35653 Le Rheu, France	
Date and reference of the validation report	2011-11-25 - Validation report 11/02	
Validation process according to EPPO Standard PM 7/98:	Yes	
Reference of the test description	0 PM7/103 not included in appendix Tigano et al. (2010). Genetic diversity of the root knot nematode Meloidogyne enterolobii and development of a scar marker for this guava-damaging species. Plant Pathology.	
Is the test the same as described in the EPPO DP?	No not included in PM7/103	
Is the lab accredited for this test?	No	
Plant species tested (if relevant)		
Matrices tested (if relevant)	Isolated nematodes	
List of methods used		
Method for extraction / isolation / baiting of target organism from matrix		
Molecular methods, e.g. hybridization, PCR and real time PCR	X	Species specific SCAR PCR
Serological methods: IF, ELISA, Direct Tissue Blot Immuno Assay		
Plating methods: selective isolation		
Bioassay methods: selective enrichment in host plants, baiting, plant test and grafting.		
Pathogenicity test		
Fingerprint methods: protein profiling, fatty acid profiling & DNA		

profiling		
Morphological and morphometrical methods intended for identification		
Biochemical methods: e.g. enzyme electrophoresis, protein profiling		
Other		
Analytical sensitivity (= limit of detection)		
What is smallest amount of target that can be detected reliably?	5 J2	
Diagnostic sensitivity		
Proportion of infected/infested samples tested positive compared to results from the standard test , see appendix 2 of PM 7/98		
Specify the standard test		
Analytical specificity		
Specificity value	100%	
Number of strains/populations of target organisms tested	4 populations of M. enterolobii	
Number of non-target organisms tested	28 nematodes populations	
Cross reacts with (specify the species)	none	
Diagnostic Specificity		
Proportion of uninfected/uninfested samples (true negatives) testing negative compared to results from a standard test		
Specify the standard test		
Reproducibility		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	100% (for 5 J2)	
Repeatability		
Provide the calculated % of agreement for a given level of the pest (see PM 7/98)	100% (for 5 J2)	
Test performance study		
Test performance study?	No	
Include brief details of the test performance study and its output.It available, provide a link to published article/report		
Other information		

Any other information considered useful e.g. robustness, ease of performing the test, etc.	Test evaluated by other teams but with mix and amplification conditions different from the article. These teams got better sensitivity.
The following complementary files are available online:	<ul style="list-style-type: none"><li data-bbox="738 297 1123 327">• Validation report ref 11/02